

ARL is an Authority on Nutrition and the Science of Balancing Body Chemistry Through Hair Tissue Mineral Analysis!

Hair Tissue Mineral Analysis

home About

Hair Analysis

Lab Profile

Educational Material

Mineral Information

Contact

Phosphorus

Home » Mineral Information » Phosphorus



Sources Of Phosphorus

| Sources O | i Pilospilorus |
|-----------------|--|
| Seafood - | tuna, mackerel, pike, red snapper, salmon, sardines, whitefish, scallops, shad, smelt, anchovies, bass, bluefish, carp, caviar, eel, halibut, herring, trout |
| Meats - | liver (beef, chicken, hog, lamb), rabbit, sweetbreads, turkey, beef brains, chicken, eggs, egg yolk, lamb heart, kidney |
| Nuts/seeds - | pinon, pistachios, pumpkin, sesame, sunflower, walnuts, almonds, brazil nuts cashews, filberts, hickory, peanuts, pecans |
| Vegetables - | chickpeas, garlic, lentils, popcorn, soybeans |
| Dairy - | cheeses |
| Grains - | wheat bran and germ, wild rice, buckwheat, millet, oats, oatmeal, brown rice, rice bran, rye, wheat |
| Miscellaneous - | chocolate, kelp, yeast, bone meal |

Roles In The Body

- Bone structure 80-85% of phosphorus in the body is located in the bones and teeth
- Energy production (ATP adenosine triphosphate and ADP adenosine diphosphate)
- Cell membranes (as phospholipids)
- Genetic reactions in DNA deoxyribonucleic acid and RNA ribonucleic acid
- Buffering agent, to maintain osmotic pressure

Functions Of Phosphorus

| Digestive - | regulates absorption of calcium and a variety of trace elements. Phosphorus in excess has a laxative action |
|------------------|---|
| Nervous - | source of adenosine triphosphate (ATP), component of the myelin sheath |
| Endocrine - | interacts with vitamin D |
| Blood - | red blood cell (RBC) metabolism |
| Muscular - | adenosine triphosphate (ATP) needed for muscle contraction |
| Skeletal - | component of bone and teeth |
| Immune - | adenosine triphosphate (ATP) for leukocytes |
| Metabolic - | energy production via phosphorylation reactions |
| Detoxification - | in liver - via adenosine triphosphate (ATP) |

Synergetic Nutrients

| Absorption - | sodium, potassium, low calcium diet, vitamin D, parathyroid hormone, high fat diet |
|--------------|--|
| Metabolic - | calcium, magnesium, B-complex vitamins (in energy production) |

Antagonistic Nutrients

| Absorption - | calcium, aluminum, iron, magnesium, vegetarian |
|--------------|--|
| | diets,vitamin D deficiency |

Hair Analysis Notes

High Hair Phosphorus:

- An elevated phosphorus level is frequently indicative of excessive protein breakdown of body tissues. As proteins break down, phosphorus is released.
- Phosphorus levels may increase temporarily as toxic metals are being eliminated in the course of a nutrition program.
 Very high phosphorus (greater than 25 mg%) can indicate a serious metabolic disturbance.
- Pubic hair samples often show elevated phosphorus readings. This is a characteristic of pubic hair.

Low Hair Phosphorus:

- A low phosphorus level is frequently associated with inadequate protein synthesis.
- Although most diets are adequate in phosphorus, those on low-protein diets or vegetarians may have a low phosphorus intake.
- Zinc is required for protein synthesis. Often a low phosphorus level is associated with a zinc deficiency, cadmium toxicity, or zinc loss. When these imbalances are corrected, the phosphorus level improves.
- A low phosphorus level may be due to poor digestion or assimilation of protein. This may be due to digestive enzyme deficiency, low hydrochloric acid level, or other factors.

This material is for educational purposes only
The preceding statements have not been evaluated by the
Food and Drug Administration
This information is not intended to diagnose, treat, cure or prevent any disease.